

## LIST OF FIGURES

1. Sites in the Albemarle-Pamlico Estuary where blue crabs were collected in this study..... p. 19
2. Sites in the Pamlico River where blue crabs were collected in this study..... p. 20
3. Effect of blue crab hemolymph on the survival of Escherichia coli D31 as measured by the plating assay. Two replicate experiments yielded similar results ..... p. 27
4. Effect of sodium chloride on the antibacterial activity of blue crab hemolymph against Escherichia coli D31 as measured by the turbidometric assay. Two replicate experiments yielded similar results ..... p. 29
5. Antibacterial activity of intermolt, clinically normal, male blue crabs collected from different geographic locations in the Albemarle-Pamlico Estuary. Values are expressed as the concentration of bacteria present after exposure to hemolymph followed by a standardized incubation period in culture medium. Thus, the lowest bacterial growth indicates the highest antibacterial activity. BR = Bay River, BS = Bogue Sound, CS = Core Sound, LR = Long Shoal River, NR = Neuse River, PA = Pamlico River, PO = outer Pamlico River (includes Rose Bay and Swanquarter Bay), PU = Pungo River. Values are expressed as predicted least square means, as described in the text. Values with (\*) are significantly different from mean activity of Core Sound crabs; values with (o) are significantly different from mean activity of Bogue Sound crabs ( $t = 1.96927$ ,  $p < 0.05$ ). No other values are significantly different from each other. Ranking of sites is from lowest (Pungo River) to highest (Bogue Sound) mean salinity based upon the hydrographic data of Wood (1967)..... p. 31
6. Antibacterial activity of intermolt male blue crabs that were either clinically normal or had clinical signs of "Pamlico River shell disease". Values are expressed as the concentration of bacteria present after exposure to hemolymph followed by a standardized incubation period in culture medium. Thus, the lowest bacterial growth indicates the highest antibacterial activity. Values are expressed as predicted least square means, as described in the text. Clinically normal crabs had significantly ( $t = 1.96927$ ,  $p < 0.05$ ) greater antibacterial activity..... p. 32